

IJDSCR : Dr.Angel Vaidic Publication Available Online at: http://www.ijdscr.org Volume – 2, Issue – 6, November - December - 2020, Page No. : 20 - 24

Effect Of Edentulism On Oral And General Health

¹Dr. Sandeep Rajan, Assistant Professor, Department of OMFS, Govt. Dental College, Alappuzha, Kerala, India. ²Dr. Ajith kumar, Assistant Professor, Department of OMFS, Govt. Dental College, Alappuzha, Kerala, India. ³Dr Naveen Nandagopal, Senior Resident, Department of OMFS, Govt. Dental College, Alappuzha, Kerala, India. ⁴Dr. Shahindah Shah, Department Of Orthodontics And Dentofacial Orthopedics Maharishi Markandeshwar College Of Dental Sciences And Research, MMU, Mullana, Haryana, India

Corresponding Author: Dr. Sandeep Rajan, Assistant Professor, Department of OMFS, Govt. Dental College, Alappuzha, Kerala, India.

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

However, there are intra- and intercountry variations in the prevalence of complete edentulism [3], and direct comparison between national samples is difficult because of the impact of various factors like education, economic circumstances, lifestyle, oral health knowledge and beliefs, and attitudes to dental care [4]. In the United States, the number of edentate individuals is likely to stay stable at 9 million and, according to the most recent information, the prevalence of edentulism amongst adults over 60 years of age was 25% [5]. In 2010, the overall rate of edentulism in Canada was 6.4%, and among adults between 60 and 79 years of age, it was 21.7% [6]. The rate of edentulism tends to vary among different regions within a country. In Canada, there is a wide variation between provinces, from 14% (Quebec) to 5% (Northwest Territories) due to associated factors such as access to fluoridated water and smoking

Studies show that edentulism is closely associated with socioeconomic factors and is more prevalent in poor populations and in women [3, 16]. In 2003, the ratio of edentulism was 6 times higher in low-income than in higher income Canadian families [3]. Other factors contributing to the

prevalence of complete tooth loss are age, education, access to dental care, dentist/population ratios, and insurance coverage [17, 18]. Most edentate people are elders who wear complete dentures in one or both jaws. Studies have demonstrated that denture wearing continues to increase due to the increase in the aging population; a large number of people still depend on removable dentures for oral function [2].

Edentulism can lead directly to impairment, functional limitation, physical, psychological, and social disability, and handicap [19]. Thus, the impact of edentulism on general health should be examined by analyzing the major dimensions of health: physical symptoms and functional capacity, social functioning and perception of well-being. This means that well-quantified endpoints of demographic significance can be used to understand the global burden of this disease. The literature has been reviewed accordingly.

Impact Oral Health

which leads to a reduction in the height of alveolar bone and the size of the denture bearing area. This reduction affects face height and facial appearance, which are altered following total tooth loss [20]. The loss of alveolar bone height and width also leads to substantial changes in the soft-tissue profile, such

Corresponding Author: Dr. Sandeep Rajan, ijdscr, Volume - 2 Issue - 6, Page No. 20 - 24

as protrusion of the mandibular lip and chin [23]. There exists an interpatient variation in these anatomic degenerative changes, and the etiology of these is still unclear. It is believed that a combination of local and systemic factors may be contributors; these include age, gender, duration of edentulism, parafunctional habits, general health, and several diseases [24].

According to a systematic review evaluating the relationship between oral function and dentition, tooth numbers below a minimum of 20 teeth, with nine to 10 pairs of contacting units, are associated with impaired masticatory efficiency, performance, and mastica- tory ability (an individual's perception of his/her ability to chew) [25].

Although some evidence suggests that reduced oral function in elders is related to muscle atrophy, aging alone has little impact on masticatory performance [29]. Most studies agree that denture wearers have only about one- fifth to one-fourth the bite strength and masticatory force of dentate individuals [30]. Furthermore, complete denture wearers require 7 times more chewing strokes than those with natural dentitions to be able to cut food into half of

its original size [31]. Moreover, the thickness of the masseter muscle was found to be decreased in edentulous patients, thus decreasing bite force [32]. This may partly explain why individuals wearing complete dentures have difficulty chewing hard foods.

This disability could substantially influence the desire to bite, to chew, and to swallow and could lead to a modification of food choices [33, 34]. As a result, research has consistently demonstrated that tooth loss and dental status have a negative impact on diet and food selection [20, 35].

Tooth Loss: Determinant of Oral Health. Edentulism can be accompanied by functional and sensory deficiencies of the oral mucosa, oral musculature, and the salivary glands. Decreased tissue regeneration and decreased tissue resistance are expected in the edentulous population, which can impair the

© 2020 IJDSCR, All Rights Reserved

protective function of the oral mucosa. Associations have been reported between aging, denture use, and oral mucosal disorders, including denture stomatitis, an inflammatory condition of the palatal mucosa seen in complete denture wearers, angular cheilitis, oral candidosis, and traumatic ulcers [36-38]. According to MacEntee et al., the odds of finding hyperplasia, stomatitis, and angular cheilitis increase approximately threefold in denture wearers [36]. Such dis- orders could expose the individual to internal and external pathogens, and their prevalence is an important parameter in evaluating the oral health of an elderly population [39, 40]. Although a direct correlation between edentulism and aspira- tion pneumonia has not been reported, the potential relation between denture plaque and aspiration pneumonia has been discussed in susceptible individuals [41, 42]. Although the majority of oral mucosal conditions in the elderly are benign, some may become malignant, especially if the protective functions of oral mucosa are decreased [37].

Denture wearers may have additional prosthetic problems as a result of soft and hard tissue damage caused by oral dyskinesia [43]. Edentulism is also associated with tardive dyskinesia, a type of dyskinesia occurring among patients chronically treated with antipsychotic drugs [44].

Impact of Edentulism on General Health

Although evidence is accumulating to support a reciprocal relationship between oral and general health [38, 60], the mechanisms linking poor general health and tooth loss are not yet clear. A purported pathway for this association involves deleterious effects of tooth loss on nutrition that, in turn, impacts systemic health [61]. Nutritional factors, especially antioxidants, may decrease following tooth loss and modulate systemic disease by interfering with the inflam- matory cascade and preventing carcinogenesis [61]. A study on 83,104 US women [45] showed that diet might partially explain the association between oral health and cardiovas- cular disease. In this cross-sectional analysis, the edentulous women had dietary

intakes associated with an increased rate of cardiovascular disease. These results are supported by a longitudinal analysis on 41,891 adults, which confirms an association between tooth loss and the prevalence of heart diseases [53].

Furthermore, excessive intakes of highly processed high- fat and high-carbohydrate foods contribute to obesity and

Furthermore, excessive intakes of highly processed high- fat and high-carbohydrate foods contribute to obesity and

in edentulous people [71]. Joshipura et al. [72] collected dietary intake data from 49,501 male health professionals and demonstrated that, compared to dentate individuals, edentu- lous respondents consumed fewer vegetables, less fiber, and less carotene intake, while consuming more cholesterol and saturated fats. These differences were independent of sociode- mographic and health behaviour characteristics. Lowe et al.

[73] established that total tooth loss was associated with low citrus fruit consumption, low plasma vitamin C levels, and increased amounts of inflammatory reactants, such as plasma C-reactive protein. They also demonstrated increased levels of plasma interleukin-6, fibrinogen, and factor VIII in women. These factors are associated with an increased risk of coronary heart diseases and stroke.

In relation to weight gain, the results of a study carried out by Lee et al. [74] demonstrated that edentulism was associated 5% in one year. Furthermore, an asso- ciation between edentulism and obesity was found in several studies [14, 75]. When edentulism was not rehabilitated with complete dentures, it was associated with both underweight and overweight/obesity in an elderly population [76].

Despite this evidence, some findings contradict the association between dentition and nutrition [77, 78]. In a cross-sectional study, Shinkai et al. [77] investigated the influence of dentition status on overall diet quality. The authors concluded that, although individuals with better dentition status had better masticatory performance and bite force, no association was found between dentition status

and quality of diet. However, in the same study, they found an association between masticatory variables and intakes of specific dietary components, such as vitamin C and fiber. There also exist some contradicting results regarding the influence of sociodemographic variables on

the dentition-nutrition relationship. Findings of Nowjack-Raymer and Sheiham [66] demonstrated that the association between dentition and nutrition was independent of age, sex, raceethnicity, and socioeconomic factors, whereas Lee et al.

[74] demonstrated racial-ethnic differences in dietary intake patterns, showing that the food intake of African American edentulous elders was similar to those with teeth. How- ever, Caucasian edentate elders displayed different dietary food patterns than their dentate counterparts. This ethnic difference could be explained by fundamental differences in socioeconomic characteristics of racial groups. African Americans consumed more fat, fewer vegetables, and less fiber than did the Caucasians, irrespective of dental condition [79].

Although diet has been shown to be poorer in edentulous populations, there is still a need for more research about the association between tooth loss and specific changes in nutrient intake. The association between tooth loss and aging

[17] may become even more important with the growth of the elder population worldwide. Such an increasing population has higher prevalence of chronic conditions [80] that may be indirectly aggravated by edentulism [73].

Regarding life expectancy of edentulous individuals, tooth loss was found to be associated with the onset of disability and mortality, even after adjusting for confounding factors such as socioeconomic and health behavior factors [81], and one study demonstrated that each tooth that remains in the oral cavity after the age of 70 decreased the risk of mortality over 7 years by 4% [82]. In addition, several studies established an association between edentulism before the age of 65 and an increased risk of earlier death [83]. Also,

© 2020 IJDSCR, All Rights Reserved

according to Shimazaki et al. [84], the mortality rate of edentulous elders without dentures was significantly higher than those with 20 or more teeth and, in a large cohort study, an association was found between tooth loss and mortality, in addition to death resulting from gastrointestinal cancer, heart disease, and stroke [46].

Impact of Edentulism on the Quality of Life

Increasingly, it is recognized that patients' perceptions of their oral health are important in evaluating well-being and determining health care outcomes [91]. The exclusive use of clinical measures has been generally criticized because they provide little insight into the psychosocial aspects of health and do not adequately reflect the health status, functioning, and perceived needs of edentulous and elderly individuals [33, 92, 93]. Edentulism may lead to changes in most of the domains leading to poorer QoL (e.g., impaired masti- cation, denture trauma, aesthetic concerns, or negative selfperception). Teeth have an important role in facial appearance, speech, and eating ability. There is overwhelming evidence showing the negative effect of edentulism on OHQoL [94, 95]. Edentulism negatively influences not only oral function, but also social life and day-to-day activities [96]. Compromised oral function has been linked to decreased selfesteem and a decline in psychosocial well-being [97]. Edentulous people may avoid participation in social activities because they are embarrassed to speak, smile, or eat in front of others, leading to isolation [98]. Many people develop skills to overcome the limitations of dentures, but some are unable to do so [20]. Fiske et al. [99] demonstrated that denture wearers have decreased self-confidence, premature aging, altered self-image, and altered behaviour in socializing and forming close relationships. On the other hand, den- tures could improve oral appearance and social interactions of individuals, which might enhance self-esteem and thus contribute to psychological well-being [90, 100]. Variables, including type of treatment, age, sex, and marital status, could explain the variation in ratings of OHQoL and tooth loss [90].

Conclusion

Edentulism has a series of deleterious consequences for oral and general health. Oral consequences vary from the well- known residual ridge resorption to an impaired masticatory function, an unhealthy diet, social disability, and poor oral health quality of life. Edentulous individuals are also in greater risk for different systemic diseases and an increase in mortality rate. Therefore, oral health care providers should prevent tooth loss with proper dental education, oral health promotion, and a high level of dental care in an attempt to assure the existence of a physiologic dentition.

References

- Health Canada, Report on the Findings of the Oral Health Component of the Canadian Health Measures Survey 2007–2009, Ministry of Health, Ottawa, Ontario, 2010.
- R. Moreira, Tooth-loss in adults and the elderly in Brazil: the influence of individual, contextual and geographical features [Ph.D. thesis], Faculdade de Saúde Pública, Universidade de Sao Paulo, Sao Paulo, Brazil, 2009.
- B. A. Dye, X. Li, and G. Thorton-Evans, "Oral health disparities as determined by selected healthy people 2020 oral health objectives for the United States, 2009-2010," NCHS Data Brief, 2012.
- M. T. Ribeiro, M. A. Rosa, R. M. Lima et al., "Edentulism and shortened dental arch in Brazilian elderly from the National Survey of Oral Health 2003," Revista de Saúde Pública, vol. 45, no. 5, pp. 817–823, 2011.
- C. E. Medina-Solís, R. Pérez-Núñez, G. Maupomé et al., "National survey on edentulism and its geographic distribution, among Mexicans 18 years of age and older (with emphasis in WHO age groups)," Journal of Oral Rehabilitation, vol. 35, no. 4, pp. 237–244, 2008.

- M. V. Eustaquio-Raga, J. M. Montiel-Company, and J. M. Almerich-Silla, "Factors associated with edentulousness in an elderly population in Valencia (Spain)," Gaceta Sanitaria, vol. 27, no. 2, pp. 123–127, 2012.
- P. Tramini, S. Montal, and J. Valcarcel, "Tooth loss and asso- ciated factors in long-term institutionalised elderly patients," Gerodontology, vol. 24, no. 4, pp. 196–203, 2007.
- B. G. Dogan and S. Gokalp, "Tooth loss and edentulism in the Turkish elderly," Archives of Gerontology and Geriatrics, vol. 54, no. 2, pp. 162–166, 2012.
- T. Ö sterberg, D. K. Dey, V. Sundh, G. E. Carlsson, J. O. Jansson, and D. Mellström, "Edentulism associated with obesity: a study of four national surveys of 16 416 Swedes aged 55–84 years," Acta Odontologica Scandinavica, vol. 68, no. 6, pp. 360–367, 2010.
- M. Madlena, P. Hermann, M. Jahn, and P. Fejerdy, "Caries prevalence and tooth loss in Hungarian adult population: results of a national survey," BMC Public Health, vol. 8, article 364, 2008.
- C. Bedos, J. M. Brodeur, L. Boucheron et al., "The dental care pathway of welfare recipients in Quebec," Social Science & Medicine, vol. 57, no. 11, pp. 2089–2099, 2003.
- H. W. Elani, S. Harper, P. J. Allison, C. Bedos, and J. S. Kaufman, "Socio-economic inequalities and oral health in Canada and the United States," Journal of Dental Research, vol. 91, no. 9, pp. 865–70, 2012.
- J. M. Brodeur, M. Benigeri, M. Olivier, and M. Payette, "Use of dental services and the percentage of persons possessing private dental insurance in Québec," Journal of the Canadian Dental Association, vol. 62, no. 1, pp. 83–90, 1996.
- 14. D. Locker, "Measuring oral health: a conceptual framework,"Community Dental Health, vol. 5, no. 1, pp. 3–18, 1988.

- 15. P. F. Allen and A. S. McMillan, "A review of the functional and psychosocial outcomes of edentulousness treated with complete replacement dentures," Journal of the Canadian Dental Association, vol. 69, no. 10, p. 662, 2003.
- 16. A. Tallgren, "The continuing reduction of the residual alveolar ridges in complete denture wearers: a mixedlongitudinal study covering 25 years," The Journal of Prosthetic Dentistry, vol. 27, no. 2, pp. 120–132, 1972.