

### Symptoms Of Dental Trauma In An Adult In A Tertiary Hospital In Pondicherry

<sup>1</sup>Dr. Shankar Narayan Sharma, Professor, Department of General Medicine, MKCG Medical College Hospital, Berhampur, Odisha, India

<sup>2</sup>Dr.Om Mishra, Junior Resident, Department of General Medicine, MKCG Medical College Hospital, Berhampur, Odisha, India

<sup>3</sup>Dr.DiptimayeeTripathy, Professor, Department of General Medicine, MKCG Medical College Hospital, Berhampur, Odisha, India

<sup>4</sup>Dr. Vishal Choudhary, Senior Lecturer, Department of Oral Medicine and Radiology, Triveni Institute of Dental Sciences, Hospital and Research Centre, Bilaspur, Chhattisgarh, India

<sup>5</sup>Dr. Gourav Grover, Post Graduate Student, Department of Oral Medicine and Radiology, Genesis Institute of Dental Sciences, Ferozepur Punjab, India.

**Citation Of This Article:** Dr. Shankar Narayan Sharma, Dr.Om Mishra, Dr.DiptimayeeTripathy, Dr. Vishal Choudhary, Dr. Gourav Grover, “Symptoms Of Dental Trauma In An Adult In A Tertiary Hospital In Pondicherry”, IJDSR May-June - 2020, Vo2. – 2, Issue -3, P. No. 20-24.

**Copyright:** © 2020 Dr. Shankar Narayan Sharma, et al. This is an open access journal and article distributed under the terms of the creative commons attribution non commercial License. This allows others to remix, tweak, and build upon the work non commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Corresponding Author:** Dr. Shankar Narayan Sharma, Professor, Department of General Medicine, MKCG Medical College Hospital, Berhampur, Odisha, India

**Type of Publication:** Original Research Article

**Conflicts of Interest:** Nil

#### Abstract

The posterior occlusal trait was assessed using the Angle’s classification while the anterior occlusal traits involved assessment of the horizontal and vertical occlusal relationships. Data was analysed using SPSS version 21.0. The analysis done were frequencies, cross-tabulations, one-way ANOVA and linear regression with statistical significance set at P=0.05

#### Results

A total of 250 patients were recruited for the study. Majority (88.8%) of the participants had Angle’s Class I

occlusion with over jet and open bite observed in 10.8% and 8.8% of the participants respectively There was statistically significant difference in the mean number of traumatised teeth with the different molar relationships (p=0.032), different horizontal occlusal relationships (p=0.049) and different vertical occlusal relationships (p=0.006), with the mean number of traumatised teeth among participants with Angle’s class III, over jet as well as open and deep bite significantly higher. There was no statistically significant relationship between the severity of injury and the various occlusal traits assessed.

## **Keywords**

occlusal traits, dental trauma, severity of injury

## **Introduction**

Its occurrence is associated with psychological, social, therapeutic and financial consequences that have a great burden on individuals and the society [1, 2]. Dental trauma has been linked to a number of predisposing/risk factors. Predisposing factors suggested include socio economic class, insufficient lip closure and protrusion of upper incisors [4, 9]. Some other occlusal traits which deviate from the ideal occlusion characteristics have been reported to predispose to traumatic dental injuries [10]. Furthermore, there is paucity of studies characterising the occlusal traits of adult patients with dental trauma. It is therefore worthy to consider the occlusal traits of adults who have dental trauma and see if there are any associations between the occlusal traits and characteristics of the dental trauma.

## **Methodology**

Study duration was August 2014 to July 2017. A convenience consecutive sampling design was employed. Data was obtained by means of a pre-tested questionnaire. Demographic characteristics (age, sex, occupation, level of education, marital status, religion and tribe) were documented. The other aspects of the questionnaire were history taking and clinical examination. A thorough history from the patient and the traumatic incident was obtained. The pre-injury data included biographic and demographic data, past medical history, time, location and mechanism of accident, occlusion and loss of consciousness. Superficial crazing or infractions may be identified with a direct light source, trans illuminating perpendicular to the long axis of the tooth from the incisal edge. Teeth were checked for any displacements, International Journal of Dental Sciences

intrusions, or luxations and noted. Also, the number of teeth affected (single vs. multiple) per participant was also recorded. For the purpose of analysis, the severity of the trauma was categorised into three in the order of severity from the most severe as follows: avulsion and crown-root fractures (III, traumatised teeth with pulpal involvement and teeth with combination injuries (II) and traumatised teeth without pulpal involvement (I). In cases where there were multiple traumatised teeth, the most severe injury was used for the categorisation. The posterior occlusal trait was assessed with the Angle's classification [14] as follows; Class I – The mesiobuccal cusp of the permanent maxillary first molar occludes in the buccal groove of the permanent mandibular first molar. Class II – The mesio buccal cusp of the permanent maxillary first molar occludes mesial to the buccal groove of the permanent mandibular first molar. Division 1 – with proclination of the maxillary incisor teeth Division 2 – with retroclination of the maxillary incisor teeth Class III – The mesiobuccal cusp of the permanent maxillary first molar occludes distal to the buccal groove of the permanent mandibular first molar.

The vertical occlusal relationship was classified into normal overbite (33-50%), deep bite (>50%) and reduced/open bite (3mm) and reverse overjet (mandibular incisor occluding ahead of the maxillary incisors). Data collected was analysed using the Statistical Package for Social Science (SPSS) version 21.0. The data was subjected to descriptive analysis in the form of frequencies, percentages and cross-tabulations. The difference in mean was determined by one-way ANOVA and the post hoc LSD was applied. Linear regression was used to determine any possible occlusal predictor of number of traumatised teeth. Statistical significance was set at  $P=0.05$ . The results were presented in form of

frequencies, percentages, pie charts, graphs, tables and cross tabulations.

## **Results**

A total of 250 patients were recruited for the study. The age of the participants ranged from 16 years to 71 years with a mean age of  $30.59 \pm 11.20$  years. There was a female preponderance (Figure 1) with male to female ratio of 1:1.08. Fall was the most prevalent (33.6%) aetiology reported among the participants. The least reported traumatised with 82.8% of the traumatised anterior teeth being maxillary anterior teeth (Figure 2). A few (10.4%) There was no statistically significant association found between single or multiple traumatised teeth and the severity of injury with the molar relationship as well as the horizontal occlusal International Journal of Dental Sciences 20 relationship. In like manner, there was also no statistically significant relationship between the severity of injury and vertical occlusal relationship among the participants.

However, there was a statistically significant association between the number of teeth affected per participant and vertical occlusal relationship among the participants ( $P=0.04$ ), with a higher proportion of participants with deep bite and open bite having multiple traumatised teeth and a higher proportion of those with normal overbite having a single tooth traumatised shows That there was Statistically significant difference in the mean number of traumatised teeth with higher than among those with Angle's class I and class II division 1. There was also statistically significant difference in the mean number of traumatised teeth with the different horizontal occlusal relationships ( $p=0.049$ ), with the mean number of traumatised teeth among participants with increased overjet significantly higher than among those with normal overjet (Figure 3). Table 5 shows that there was statistically significant difference in the mean number of

aetiology was iatrogenic dental event accounting for 0.8% (Table 1). Majority (88.8%) of the participants had Angle's Class I occlusion with over jet and open bite observed in 10.8% and 8.8% of the participants respectively (Table 2). The participants presented with a range of 1 to 9 traumatised anterior teeth with the mean number of traumatised teeth being  $1.88 \pm 1.28$ . About half (49.6%) of the participants had only a single tooth.

traumatised teeth with the different vertical occlusal relationship ( $p=0.006$ ), with the mean number of traumatised teeth among participants with open and deep bite significantly higher than among those with normal bite.

## **Discussion**

Trauma to the anterior teeth is a common occurrence and although, occurrences are high in childhood and adolescence, the incidence in adulthood peaks at 21-30 years [3]. Many studies have been carried out on children to establish the relationship between trauma and certain occlusal traits. This study was designed to assess the relationship between anterior and posterior occlusal traits and characteristics of trauma to the anterior teeth among adult patients who reported to the clinic for reasons other than orthodontic and were diagnosed with one form of traumatic dental injury or the other. The molar, horizontal and vertical occlusal relationships were assessed. The injury pattern in this study showed that there were more complex injuries involving the pulp and avulsion. This is in consonant with previous findings where adults were observed to have more complex injuries [15]. This maybe because local biological factors may affect injury pattern [15] as the mechanical properties of the teeth and bone differ in adulthood.16-18 Although, Angle's class II with

increased overjet has been reported as a predisposing factor for dental trauma [3, 19, 21], it was observed in this study that most of the participants had Angle's class I molar relationship with normal overjet and overbite. A finding which is different from that reported in a previous study where a greater number of patients with trauma were found to have class II division 1 malocclusion [22]. This may be due to the high prevalence of Angle's class I malocclusion and normal overjet in our environment [23, 24]. More so, most of the previous studies were done on children and adolescents [19, 20] who are more prone to trauma and traumatic injuries. Also, the aetiology of dental trauma seems to differ between children and adults [15]. Various occlusal traits have been associated with increased dental trauma. Shulman and Peterson [3] observed that the odds of trauma increased with increased overjet while Antunes et al [1] reported that increased overjet was three times more likely to be associated with trauma than normal overjet. Overjet can be said to be a predisposing factor for trauma and was found not to influence the severity of the trauma in this study. This may be because the extent of damage to teeth following dental trauma tends to be determined by the energy and direction of the impact as well as the resilience of the involved structures [15, 25]. The mean number of traumatised teeth (1.88) recorded in this study was lower than that reported by Teshome and Muche (2.31) [26] as well as that by Ruslin (3.55) [27]. In contrast the prevalence of single traumatised tooth (49.6%) observed in this study was noted to be higher than that reported in previous studies [26, 28, 29]. The mean number of traumatised teeth was observed to be associated with molar relationship with the mean number of traumatised teeth among participants with Angle's class III significantly higher. Angle's Class III malocclusion tends

to be associated with dental compensations in which the maxillary incisors are excessively flared and the mandibular incisors severely tipped [30] probably making them more prone to trauma. The mean number of traumatised teeth among participants with increased overjet was observed to be significantly higher. Artun et al [31] found a strong association between horizontal occlusal relationship and increasing trauma occurrence. Shulman and Peterson [3] reported that after adjusting for age, overjet was the only occlusal covariate significantly associated with maxillary incisor trauma, with the odds of trauma increasing markedly as overjet increased. It can be inferred based on this finding that overjet is not just a predisposing factor but can influence the mean number of teeth involved in a traumatic injury. This can be attributed to the fact that increased overjet means increased exposure of teeth with inadequate lip protection which predisposes more teeth to trauma [26]. There appears to be contrasting reports in the literature on the association between dental trauma and vertical occlusal relationships [31, 34]. While Bonini et al [32] and Oliveira et al [33] reported significant relationships between anterior open bite and dental trauma in their studies, Goettems et al [34] and Artun et al [31] in their studies could not establish any relationship. However, Goettems et al [34] were able to establish a relationship between deep bite and dental trauma. Interestingly, a strong link between the vertical occlusal relationship and the mean number of teeth traumatised per participant was noted in this study with the mean number of traumatised teeth significantly higher among participants with open and deep bite. Further studies with larger study population assessing more occlusal traits and orthodontic parameters be done.

## References

1. Sgan-Cohen HD, Megnagi G, Jacobi Y, Dental trauma and its association with anatomic, behavioural and social variables among fifth and sixth grade school children in Jerusalem. *Community Dent Oral Epidemiol.* 2005; 33:174- 180.
2. Aldrigui JM, Abanto J, Carvalho TS, Mendes FM, Wanderley MT, Bonecker M. et al. Impact of traumatic dental injuries and malocclusion on quality of life of young children. *Health Qual Life Outcomes.* 2011; 9:78-86
3. Al-Sharheed M, Bedi R, Hunt NP. Traumatized permanent teeth in 11 to 16-year-old Saudi Arabian children with a sensory impairment attending special schools. *Dent Traumatol.* 2003; 19:123-125. 1
4. Glendor U. Aetiology and risk factors related to traumatic dental injuries: A review of the literature. *Dent Traumatol.* 2009; 25:19-31.
5. Otuyemi OD. Traumatic anterior dental injuries related to incisor overjet and lip competence in 12-year-old Nigerian children. *International Journal of Paediatric Dentistry.* 1994; 4:81-85.
6. Adekoya-Sofowora CA. Traumatized anterior teeth in children: a review of the literature. *Niger J Med.* 2001; 10(4):151-7.
7. El-Karmi RF, Hamdan MA, Rajab LD, Abu-Ghazaleh SB, Sonbol HN. Prevalence of traumatic dental injuries and associated factors among preschool children in Amman, Jordan. *Dent Traumatol.* 2015; 31:487-492.
8. Angle EH. Classification of malocclusion. *Dent Cosmos.* 1899; 4:248-64.
9. Lauridsen E, Hermann NV, Gerds TA, Andreasen JO. Pattern of traumatic dental injuries in the permanent dentition among children, adolescents and adults. *Dent Traumatol.* 2012; 28:358-363.
10. Antunes LA, Leão AT, Maia LC. The impact of dental trauma on quality of life of children and adolescents: A critical review and measurement instruments. *Cien Saude Colet.* 2012; 17:3417-24.
11. Arhakis A, Athanasiadou E, Vlachou C. Social and psychological aspects of dental trauma, behaviour management of young patients who have suffered dental trauma. *Open Dent J.* 2017; 11:41-47.