

Dental Diseases Among Children Attending Dental Department In Kolkata Child Hospital, Kolkata, India

¹Dr. Ram Prakash Mishra, Post Graduate 3rd Year, ITS Dental College, Hospital And Research Center, Knowledge Park III, Uttar Pradesh, India

²Dr. Bhavana Kumari, Postgraduate 2nd Year, Department Of Oral And Maxillofacial Surgery, ITS Dental College, Hospital And Research Center, Uttar Pradesh, India

³Dr. Prashant Kumar, Postgraduate 2nd Year, Department Of Oral And Maxillofacial Surgery, ITS Dental College, Hospital And Research Center, Uttar Pradesh, India

⁴Dr. Radhika Devi, Principal, Department Of Oral And Maxillofacial Surgery, ITS Dental College, Hospital And Research Center, Thiruvananthapuram, Kerala, India

Citation Of This Article: Dr. Ram Prakash Mishra, Dr. Bhavana Kumari, Dr. Prashant Kumar, Dr. Radhika Devi, “Dental Diseases Among Children Attending Dental Department In Kolkata Child Hospital, Kolkata, India”, IJDSQR May-June - 2020, Vo2. – 2, Issue -3, P. No. 29-32.

Copyright: © 2020 Dr. Prashant Kumar, et al. This is an open access journal and article distributed under the terms of the creative commons attribution non commercial License. This allows others to remix, tweak, and build upon the work none commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Corresponding Author: Dr. Prashant Kumar, Postgraduate 2nd Year, Department Of Oral and Maxillofacial Surgery, ITS Dental College, Hospital and Research Center, Uttar Pradesh, India

Type of Publication: Original Research Article

Conflicts of Interest: Nil

Abstract

Oral and dental diseases such as dental caries and its sequelae, periodontal diseases are still major public health problems worldwide, but more in developing countries. In context with Bangladesh due to lack of research and evidence in this field , a baseline data regarding the occurrence of oral diseases has been seen to be necessary in order to plan any community oriented preventive programs.

Objective

To find out Pattern of oral and dental diseases among children attending Dental Department in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh.

Materials and Methods

This was a cross sectional hospital based study carryout among 443 children (242 males and 201 females) attending in the Department of Pediatric Dentistry in Dhaka Shishu (Children) Hospital. All the new patients attending to dental OPD with the complained of oral and dental problems were included in this study. Old patients and those who were not willing to participate in the clinical examination for study were excluded from the study. The study period was June 2019 to August 2019.

Conclusion

Dental caries and its sequelae was the most predominant disease in the children followed by oral mucosal diseases. Oral health promotion and planning programs directed

towards the oral health should be conducted at regular intervals.

Keywords

children, oral and dental diseases; cross-sectional study.

Introduction

Oral and dental health is the standard of the oral and related tissue, which enables an individual to eat, speak, and socialize without active discomfort and embarrassment which contributes to general well-being of the individual [1]. Oral and dental diseases are common in the developing countries. The pattern of these diseases vary from one community to another, and depends on various factors like oral health care practice the level of dental awareness. With sufficient manpower, availability of modern dental equipment and a wide range of dental materials, oral health care for the child has improved in the past three decades. This has been attributed to advances in technology [4].

During this visit the developing occlusion should be monitored throughout eruption at regular clinical intervals [6]. It has been shown that pre-school children who visited the oral and dental health center early will more likely visit for preventive reasons as adults [7]. Epidemiological studies have shown that the prevalence of dental diseases particularly dental caries is on the steady increase in developing countries as refined sugar, sweets and fruit drinks have become more available [8-11]. Currently dental caries has become a serious problem in school children and periodontal disease is wide spread in adult population [12]. The purpose of the study is to assess the pattern of distribution of oral and dental diseases among children aged 1 month to 14 years visiting the hospital within a 3 months' time period.

To find out the Pattern of oral and dental diseases among children attending Dental Department in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh.

Materials and Methods

This was a cross sectional hospital based study carried out among 443 children (242 males and 201 females) attending in the Department of Pediatric dentistry in Dhaka Shishu (Children) Hospital, Dhaka, Bangladesh. All the new patients attending to dental OPD with the complaint of oral and dental problems were included in this study. Old patients and those who were not willing to participate in the clinical examination for study were excluded from the study. The study period was June 2019 to August 2019. All the patients were examined clinically and investigations were carried out for the required cases. The demographic data, pattern of oral and dental diseases were recorded. Data were analyzed by chi-squared test by using SPSS Windows version 22.0.

Results

The study was carried out among (443) children patients who were attending in the dental OPD of Dhaka Shishu (Children) Hospital during the period of June 2019 to August 2019. There were 242 (54.6%) males and 201 (45.4%) females patients, a male-female ratio of 1.2:1. The age range of children were 224 (50.6%) below the age of 5 years, 200 (45.1%) at the age of 5-10 years and 19 (4.3%) above 10 years (10-14 years) are shown in [Table 1 to 2]. [Table-1] shows the highest number of child patient's cases in below 5 years 224 (50.6%) and lowest above 10 years out of 443 (7.3%) patients. Shows highest number of patients were male [Table-2].

The total Association of dental disease with different age. Among them Caries disease all group average high rate so P-Value (0.010*). [Table-6] So grossly destructed due to caries 49 cases of <5 yrs. 35.5%, 85 cases of 5-10yrs. 54.1%, and 7 cases of under >10 yrs. 5.1%. Significant and

all patients are another case disease. It was also seen that the prevalence of dental caries was somewhat similar among males (80.5%) and females (80.8%). But the younger children aged between 5 to 10 years showed more dental caries (85.5%) than the older children of above 10 to years 6 (3.8%). However, this difference was not statistically significant ($p = 0.11$).

Discussion

This study was conducted to evaluate the pattern of distribution of oral and dental diseases in children in Dhaka Shishu (Children) Hospital. In this study, the predominant oral disease was found to be dental caries in 22.7% and its sequelae like grossly carious teeth 19.9%, dentoalveolar abscess-11.5% BDR 7.7% and pulpitis 7.6%. The next prevalence of oral diseases is oral mucosal disease like stomatitis- 15.7%, geographical tongue 3.5% and oral thrush-3.5%, Gingivitis 1.0%, Traumatic injury of tooth, soft tissue and bony tissue 1.0%, Pericoronitis 0.5%, Natal teeth 0.1% and other diseases 0.5%. Dental caries attributes to toothache and eventually children have to miss their school which will lead to poor performance of children in school. Severe dental decay was the main dental factor related to school absence in Thai children [13, 6]. It was also seen that the prevalence of dental caries and its sequelae like grossly carious teeth, pulpitis, dentoalveolar abscess, (retained broken root was more in the age group of 5 to 10 years old child than the age group below 5 years. Dental caries is also known to be commonest cause of tooth loss because most patients presents with advanced dental decay for which restorative treatment will not be possible. Complications arising from late presentation include dental abscess, grossly broken shown crown, retained broken roots and osteomyelitis in severe cases. It was also seen that the prevalence of dental caries was somewhat similar among males (54.6%) and females (45.4%).

But the younger children aged between 5 to 10 years showed more dental caries (85.7%) than the older children above of 10 years 6 (3.8%). Out of 443 children 157 (22.7%) showed presence of dental caries, 138 (19.9%) had grossly destructed carious teeth, 79 (11.5%) had dentoalveolar abscess, 53 (7.7%) had retained broken root, 52(7.6%) were suffering from pulpitis, 108 (15.7%) of the children presented with stomatitis, 34 (4.8%) had retained deciduous teeth, 24(3.5%) geographical tongue , 24(3.5%) oral thrush, 7(1.0%) gingivitis 7(1.0%) presented with traumatic injury of tooth, soft tissue and bony tissue, 4(0.5%) had pericoronitis, 1(0.1%) had natal tooth and 4(0.5%) were others like extra oral swelling, aphthous ulcer , mucocele. The total Association of dental disease with different age group. Among them Caries disease all group average high rate so p -Value (0.010*). So grossly destructed due to caries 49 cases below of 5 yrs 35.5%, 82 (59.7%) cases of 5-10yrs, and 7(5.1%) cases above 10 yrs. significant .However, this difference was not statistically significant ($p = 0.11$). This is in accordance with the prevalence studies in Nepal which showed that younger children presented with more dental caries than older children. Their study have reported that 5-6 yrs. old had caries 52% and 69% whereas 12-13 years old children had 41% and 53.23% caries prevalence respectively in Nepalese population [3,7]. 37(46.8%) of children aged <5 years and 49.4% of the children aged 5-10 years presented with dent alveolar abscess. This difference was found to be statistically significant ($p < 0.05$). They have also reported that the untreated dental caries was the most common cause of dent alveolar abscess in children, followed by trauma, failed restoration or periodontal infection [16]. Traumatic injuries in primary teeth is commonly more in children aged between 13 and 24 months [17].

Conclusion

In this study dental caries and its sequelae was seen to be the most prevalent oral disease among children of 1 month to 14 years of age. Therefore, it can be derived from this study should be given to the treatment and control of dental caries so as to decrease the prevalence of oral diseases as a whole. Rehabilitative care targeted towards treatment of the existing problems of dental caries should be made available to the children. Along with this, oral health education programs at individual, and community levels should be implemented aimed primarily at increasing the knowledge and awareness of the communities towards the risk factors associated with oral problems in order to prevent further increase of occurrence of dental caries among children. A significant 95.4% of the children made symptomatic visit. Oral mucosal disease was the second prevalent finding followed by dental caries. There is an urgent need to increase dental health awareness among children through Hospital based continuing dental tips and also among other pediatric care givers such as parents and pediatricians.

References

1. Enwonwu CO. Socio-economic factors in dental caries prevalence and frequency in Nigerians: an epidemiological study. *Caries Res.* 1974; 8:155-71.
2. Henshaw NE, Adenubi JO. The increase in dental disease in the Northern States of Nigeria and its manpower implications. *J Dent.* 1975; 3:243-56.
3. Jeboda SO, Ogunbodede EO. Dental carids in Africa. *Nig Qt J HospMed.* 1994; 5:57-62.
4. Noar J, Portnoy S. Dental status of children in a primary and secondary school in Zambia. *Int Dent J.* 1991; 41:142-8.
5. Ana JR. Dental manpower need in a developing community: a critical analysis of the West African scene. *Int Dent ll.* 1976; 26:411-20.

6. Krisdapong S, Prasertsom P, Rattananangsim K, SheihaA. School absence due to toothache associated with socio- demographic factors, dental caries status, and oral health related quality of life in 12- and 15-year-old Thai children. *J Public Health. Dent,* 2013 Aug 19. doi: 10.1111/jphd. 12030.
7. Prasai Dixit, et al.: Dental caries prevalence, oral and healthknowledge and practice among indigenous Chepang school children of Nepal. *BMC Oral Health* 2013.13:20. Azodo CC, Chukwumah NM, Ezeja EB. Dentoalveolar abscess among children attending a dental clinic in Nigeria. *Odontostomatol Trop.* 2012; 35(139):41-6.
8. Costa VPP, et al. "Traumatic dental injuries in primary teeth: severity and related factors observed at a specialist treatment centre in Brazil." *European Archives of Paediatric Dentistry,* 2013, 1-6.
9. Azodo CC, Chukwumah NM, Ezeja EB. Dentoalveolar abscess among children attending a dental clinic in Nigeria. *Odontostomatol Trop.* 2012; 35(139):41-6.