

## **Peripheral Glycerol Injection For Relief In Trigeminal Neuralgia And Other Neuralgic Pain Treatment**

<sup>1</sup>Dr. Karishma Pramanik, Senior Lecturer, Department of Prosthodontics, Crown, Bridge & Implantology, AME'S Dental College, Raichur, Karnataka

<sup>2</sup>Dr. Ajay Singh, Professor & HOD, Department of Prosthodontics, Crown, Bridge & Implantology, HKES's S. Nijalingappa Institute of Dental Science and Research, Kalaburagi, Karnataka

<sup>3</sup>Dr. Vandana Pareek, Post Graduate Student, Department of Prosthodontics, Crown, Bridge & Implantology, HKES's S. Nijalingappa Institute of Dental Science and Research, Kalaburagi, Karnataka

<sup>4</sup>Dr. Rajkumar Thakur, UG Student, Krishna Institute of Medical Sciences, School of Dental Sciences, Malkhapur, Karad, Maharashtra, India

<sup>5</sup>Dr. Sanjay Yadav, UG Student, Krishna Institute of Medical Sciences, School of Dental Sciences, Malkhapur, Karad, Maharashtra, India

**Citation Of This Article:** Dr. Karishma Pramanik, Dr. Ajay Singh, Dr. Vandana Pareek, Dr. Rajkumar Thakur, "Peripheral Glycerol Injection For Relief In Trigeminal Neuralgia And Other Neuralgic Pain Treatment", IJDSQR September - October - 2020, Vo2. – 2, Issue -5, P. No. 50-53.

**Copyright:** © 2020 Dr. Vandana Pareek, et al. This is an open access journal and article distributed under the terms of the creative commons attribution non commercial License. This allows others to remix, tweak, and build upon the work none commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Corresponding Author:** Dr. Vandana Pareek, Post Graduate Student, Department of Prosthodontics, Crown, Bridge & Implantology, HKES's S. Nijalingappa Institute of Dental Science and Research, Kalaburagi, Karnataka

**Type of Publication:** A Review Article

**Conflicts of Interest:** Nil

### **Abstract**

#### **Material and Methods**

Fifty patients were randomly recruited as known cases of trigeminal neuralgia of either sex and were administered glycerol injection in the peripheral branches of trigeminal nerve. Patients were followed up for next 1 year at monthly intervals and intensity of their pain was assessed. Data was recorded in specially made pro forma and analyzed using SPSS 20.0. Analysis included frequencies, mean  $\pm$  standard deviation (SD) and Paired t test. P-value

#### **Results**

Results have established positive impact of administering glycerol injection in achieving better analgesic control as compared to conventional methods. The ease in performing this procedure, early return of lost sensations and faster onset of pain relief with minimal complications makes it a worthwhile choice.

#### **Keywords**

glycerol injection, trigeminal neuralgia.

## **Introduction**

Trigeminal neuralgia characteristically affects individuals older than 40 years of age (the average age at onset is 50 years), although it may affect persons as early as puberty. Women are affected slightly more often than men, and the right side is involved more often than the left. Any branch of the trigeminal nerve may be involved, but the ophthalmic division is affected in only 5% of cases. More than one branch may be involved, and the pain is occasionally bilateral.

Spontaneous remissions occur, often lasting more than 6 months, especially during the early phase of the disease. However, the failure of medical treatment has led to the development of various surgical techniques, which include peripheral injections of various agents having neurolytic properties, [2-6] peripheral neurectomy [7] cryotherapy, microvascular decompression, radiofrequency thermocoagulation and gamma knife radiosurgery. An ideal treatment is one that causes no morbidity and preserves the normal sensation of the face. Such a sensation-preserving, absolutely safe, and permanently successful treatment does not yet exist. In some studies, more than 50% of patients with TN eventually had some kind of surgical procedure [12, 13]. Compared with other operative treatments, the peripheral injection techniques are simple, quick, and easy to perform, and they can be used as outpatient procedures.

## **Material and Methods**

The diagnosis of ITN will be made on the basis of the history and diagnostic criteria for TN. No patients with a history of any other neurologic diseases, and no evidence of organic disturbances of the face, maxillary sinus, mandible, or Temporomandibular joint will be found on radiographic examination. All patients will be treated without systemic sedation or analgesia, and all will

tolerate the procedure extremely well. Before the glycerol injection, the nerve will be carefully anesthetized and the absence of pain will be tested by sensory stimulation of the trigger zone. Ten minutes after local anesthesia, pure glycerol, which will be sterilized for 1 hour at 150°C, will be injected through a short 22-gauge needle. Injection volumes will be 0.5, 1, and 1.5 mL for infraorbital, mental, and mandibular nerve injections, respectively. All patients will be discharged with their families immediately after the injection. Patients will be advised to continue a therapeutic level of their antineuralgia medications until they will be free of pain for a full week before beginning to taper medication usage. All cases will be critically evaluated and followed for 3 years. Pain-free periods will be measured at varying intervals, including the last examination, and patients will be questioned about sensation in the injected region. All the acquired data will be then entered in SPSS 20 for data processing. Analysis included frequencies, mean  $\pm$  standard deviation (SD) and Paired t test. A p-value.

## **Results**

A total of 50 patients with trigeminal neuralgia were included in this study. There were 33 females and 17 male patients. Twenty five patients received peripheral glycerol injection (study group) and normal saline was administered to the rest of the 25 patients (control group). The right side of the face was involved in 33 cases and the left side in rest of the 17 patients. No case presented with bilateral involvement. Mandibular division was involved in 38 cases and maxillary division in 12. The control group obtained no pain relief at the next review followups. These patients continued to experience no pain relief at follow up visits. In the study group 22 patients had complete pain relief for 3 month duration. Another 7 had occasional pain during this period, with no need for

medication. One patient experienced no pain relief after the glycerol injection. After six months 14 of our patients had complete pain relief and this figure dropped to 8 at the one year followup. Overall 12 patients had no pain or occasional pain after one year, and did not require any medical therapy.

### Discussion

He also pointed out that no signs of nerve degeneration or other morphologic changes were observed in any of the experimental and control specimens at any of the time intervals studied. In one of our patients who had persistent pain despite the glycerol injection, a mental neurectomy was performed 1 week after the injection, and microscopic examination of the excised nerve confirmed the findings in the experimental study by AlKhateeb. The nerve presented inflammatory alterations only. However, it may be possible that the nerve was not accurately injected. In this respect, it would be wrong to correlate the findings of our study with that of Al-Khateeb or others. Despite high rates of initial success with virtually all surgical procedures, most doctors and patients choose medical therapy first because of potential surgical morbidity, the risk for loss of facial sensation after surgery, or the recurrence of pain despite initial surgical success. All the patients in our study had received medical treatment before the procedure, and none of them had undergone any other treatment. Medical treatment is the first choice in the management protocol for ITN patients in our clinic, which is followed by PGI, repeated PGI, neurectomy, repeated neurectomy, and scar revision at periphery of the nerve, in sequence. The current trend in surgical treatment of ITN is to use less invasive procedures. Alcohol injection is not generally preferred because it can cause peripheral necrosis. Percutaneous retrogasserian glycerol rhizotomy (PRGR) is a good example of this trend,

because pain relief can last more than 6 years with little sensory loss. Its success rate is reportedly between 82%20 and 96%2. Mild hypoesthesia, which has been reported in between 17%21 and 67%20 of patients, is apparently the only rare longterm complication of PRGR. It can be concluded that the mild hypoesthesia noted after both PGI and PRGR is the result of the effect of glycerol on the peripheral nerve, irrespective of the site of application. Most transient major complications of PRGR were not observed in the patients treated in our study. Although the onset of pain relief with PGI was immediate, the onset of pain relief after PRGR.

### Conclusion

Results concluded from data of this study shows that when the high mortality and morbidity of the other surgical procedures are compared to those associated with glycerol injection, it appears that this procedure may present a good treatment choice. It is relatively simple to perform and has a faster onset of pain relief and less complications than the other surgical treatments. However, long follow-up periods are required.

### References

1. Maness WL, Benjamin M, Podoloff R, Bobick A, Golden RF. Computerized occlusal analysis: a new technology. *Quintessence Int* 1987; 18:287-92.
2. Kerstein R, Lowe M, Harty M, Radke J. A force reproduction analysis of two recording sensors of a computerized occlusal analysis system. *J Craniomandib Pract.* 2006;24:1-10
3. Harvey WL, Osborne JW, Hatch RA. A preliminary test of the replicability of a computerized occlusal analysis system. *J Prosthet Dent* 1992;67: 697-700.
4. Hsu M, Palla S, Gallo LM. Sensitivity and reliability of the T-scan system for occlusal analysis. *J Craniomandib Disord* 1992;6:17-23

5. Mizui M, Nabeshima F, Tosa J, Tanaka M, Kawazoe T. Quantitative analysis of occlusal balance in intercuspal position using the T-Scan system. *Int J Prosthodont*. 1994 Jan-Feb;7(1):62-71.
6. A study on display and accuracy of occlusal contacts by means of T-Scan System Kanagawa Shigaku. 1990 Sep;25(2):236-41
7. Hirano S, Okuma K, Hayakawa I. In vitro study on accuracy and repeatability of the T Scan H system Kokubyo Gakkai Zasshi. 2002 Sep;69(3):194-201.
8. Koos B, Godt A, Schille C, Goz G. Precision of an instrumentation-based method of analyzing occlusion and its resulting distribution of forces in the dental arch; *J Orafac Orthop* 2010; 71:403-10.
9. Throckmorton GS.; Rasmussen J; Caloss R. Calibration of T-Scan sensors for recording bite forces in denture patients. *Journal of Oral Rehabilitation* 2009 36;636-643.
10. A study on display and accuracy of occlusal contacts by means of T-Scan System Kanagawa Shigaku. 1990 Sep; 25(2):236-41.
11. Archer S. and Vig, P.: Effects of Head Posture on Infra Oral Pressure in Classes I, II and III Adults, *Am J Orthod* 1985; 87:311- 318.
12. Ell edge DA, Schorr BL. A provisional and new crown to fit into the clasp of an existing removable partial denture. *J Prosthet Dent* 1990;63:541 – 4
13. Helvey GA. Retrofitting crowns to an existing removable partial denture clasp: a simple technique. *J Prosthet Dent* 2002; 87: 399-402.
14. Seo JM, Ahn SG. Fabrication of a metal-ceramic crown to fit an existing partial removable dental prosthesis using ceramic pressed to metal technique: a clinical report. *J Adv Prosthodont* 2014;6:241-4
15. Yoon TH, Chang WG. The fabrication of a CAD/CAM ceramic crown to fit an existing partial removable dental prosthesis: A clinical report. *J Prosthet Dent* 2012; 108;3:143-6
16. Hansen CA, Russell MM. making a crown to fit accurately under an existing removable partial denture clasp assembly. *J Prosthet Dent* 1994; 71: 206 – 8